

# What is Depression?

Symptoms of Depressive Episode			
depressed mood	loss of interest or pleasure	significant weight change	diminished concentration
sleep difficulties	fatigue nearly every day	feeling of worthlessness	recurring thoughts of death
Symptoms must cause significant distress.		Symptoms must last for at least two weeks.	

## Demographics

- Women are 2x more likely to develop depression.
- About 1 in 10 people will experience depression during their lifetime.
- Most people experience their first depressive episode between ages 20 and 30

## Risk for Depression

- Family history of depression or similar disorder
- Poverty, unemployment, social isolation, and other stressful life events
- Regular drug and alcohol use

## Psychotherapy

(Cognitive Behavioral Therapy)

CBT works by changing self-defeating thoughts and behaviours.

CBT has been found to be equally, if not more effective than medicine in many cases.

CBT is the most researched form of psychotherapy for depression.

## Medication

(Selective Serotonin Reuptake Inhibitors)

SSRIs increase the level of serotonin (a chemical related to depression) in the brain.

Studies suggest that SSRIs are the most effective when used to treat severe depression.

SSRIs don't work overnight – it might take up to 6 weeks before they reach their full effect.



A combination of both psychotherapy and medication has been found to be the most effective treatment for depression

## Other Facts

- Over ½ of those diagnosed with depression also suffer from anxiety.
- 60% of those who die by suicide suffer from depression or a related mood disorder.
- Physical exercise has been found to have a significant antidepressant effect.
- Depressive episodes also occur during bipolar disorder alongside manic episodes.

Reference: 2015 Therapist Aid LLC; Provided by **TherapistAid.com**